Mon. Not. R. Astron. Soc. 418, 2032–2042 (2011)

Ionized outflows in SDSS type 2 quasars at $z \sim 0.3-0.6^*$

M. Villar-Martín,^{1,2}† A. Humphrey,³ R. González Delgado,¹ L. Colina² and S. Arribas²

¹Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain
²Centro de Astrobiología (INTA-CSIC), Carretera de Ajalvir, km 4, 28850 Torrejón de Ardoz, Madrid, Spain
³Instituto Nacional de Astrofísica, Optica y Electrónica (INAOE), Aptdo. Postal 51 y 216, 72000 Puebla, Mexico

Accepted 2011 August 11. Received 2011 August 10; in original form 2011 July 8

ABSTRACT

We have analysed the spatially integrated kinematic properties of the ionized gas within the inner $r \leq$ few kpc in 13 optically selected SDSS type 2 quasars at $z \sim 0.3$ –0.6, using the [O III] $\lambda\lambda$ 4959,5007 lines. The line profiles show a significant asymmetry in 11 objects. There is a clear preference for blue asymmetries, which are found in 9/13 quasars at 10 per cent intensity level. In coherence with studies on other types of active and non active galaxies, we propose that the asymmetries are produced by outflows where differential dust extinction is at work.

This scenario is favoured by other results we find: in addition to quiescent ambient gas, whose kinematic properties are consistent with gravitational motions, we have discovered highly perturbed gas in all objects. This gas emits very broad lines ($R = FWHM[O III]/FWHM_{stars} \gtrsim$ 2). While the quiescent gas shows small or null velocity shifts relative to the systemic velocity, the highly perturbed gas trends to show larger shifts which, moreover, are blueshifts in general. Within a given object, the most perturbed gas trends to have the largest blueshift as well. All together support that the perturbed gas, which is responsible for the blue asymmetries of the line profiles, is outflowing. The outflowing gas is located within the quasar ionization cones, in the narrow line region.

The relative contribution of the outflowing gas to the total [O III] line flux varies from object to object in the range $\sim 10-70$ per cent. An anticorrelation is found such that, the more perturbed the outflowing gas is, the lower its relative contribution is to the total [O III] flux. This suggests that outflows with more perturbed kinematics involve a smaller fraction of the total mass of ionized gas.

Although some bias affects the sample, we argue that ionized gas outflows are a common phenomenon in optically selected type 2 quasars at $0.3 \leq z \leq 0.6$.

Key words: galaxies: ISM – quasars: emission lines – quasars: general.

1 INTRODUCTION

Evidence for an intimate connection between supermassive black hole (SMBH) growth and evolution of galaxies is nowadays compelling. Not only have SMBHs been found in many galaxies with a bulge component, but tight correlations exist between the black hole mass and some bulge properties, such as the stellar mass and velocity dispersion σ (Ferrarese & Merritt 2000; Gebhart et al. 2000). The origin of this relation is still an open question, but

†E-mail: villarmm@cab.inta-csic.es

quasar induced outflows might play a critical role. Hydrodynamical simulations show that the energy input from quasars can regulate the growth and activity of black holes and their host galaxies (Di Matteo, Springel & Hernquist 2005). Such models show that a merger between two galaxies with SMBH leads to strong inflow that feeds gas to the SMBH activating the quasar phase. The energy released by the strong outflows associated with major phases of accretion expels enough gas to quench both star formation and further black hole growth. This determines the lifetime of the quasar phase and explains the relationship between the black hole mass and the stellar velocity dispersion.

This scenario, although very attractive, still lacks compelling observational support. The presence of outflows in different classes of active galaxies has been known for several decades. However, it is

^{*}Based on observations carried out at the European Southern Observatory (Paranal, Chile) with FORS2 on VLT-UT1 (programmes 081.B-0129 and 083.B-0381).

still not clear whether they can generate important feedback effects with a notable impact on the systems evolution. The limitations are several, such as the uncertainty on where the outflows originate and their geometry, or the fact that several gas phases (molecular, neutral, ionized) can be involved. These are often observable at different spectral ranges and spatial scales and therefore require different observing techniques.

The presence of outflows of ionized gas in different types of active galaxies is well established. They are evident, for instance, as absorption lines in the ultraviolet and X-ray spectra, blueshifted relative to the systemic velocity of the host galaxy. Outflowing velocities of up to $\sim 2000 \text{ km s}^{-1}$ have been measured (see Crenshaw, Karemer & George 1999, for a review).

Outflows of ionized gas have also been discovered when characterizing the gas kinematics using the spectral properties of a diversity of forbidden lines in the infrared (e.g. Spoon & Holt 2010) and the optical. The $[O III]\lambda\lambda4959,5007$ lines have been used more frequently in optical studies. These are often the strongest optical lines and have the advantage of being cleanly isolated from other emission and absorption features. In addition, underlying contamination by line emission from the broad line region (BLR) is not expected, as is often the case for permitted lines such as H α and H β .

The predominance of blue asymmetries on the [O III] line profiles, their frequent blueshift relative to different indicators of the systemic redshift and the kinematic substructure of the lines indicate that the narrow line region (NLR) gas is not a stable, quiescent gas reservoir that follows the bulge gravitational potential. This dynamic behaviour has often been interpreted in terms of outflows where differential reddening is presen, i.e. at least part of the NLR gas is moving radially outwards and dust reddening is more severe for the receding gas. This applies to active galaxies in general (e.g. Greene & Ho 2005) and specific active galactic nucleus (AGN) types: Seyferts 1 and 2s (Heckman et al. 1981; Whittle et al. 1988; Veilleux 1991; Colbert et al 1996; Crenshaw et al. 2010), Narrow Line Seyfert 1 galaxies (Bian, Yuan & Zhao 2005; Komossa et al. 2008), radio loud and radio quiet type 1 quasars (Leipsky & Bennert 2006; Heckman, Miley & Green 1984; Boroson 2005) and narrow line radio galaxies at different redshifts (Heckman et al. 1981; Humphrey et al. 2006; Holt, Tadhunter & Morganti 2008). Heckman et al. (1981) found that blue asymmetries are in general not found in steep spectrum (extended) radio sources, but preferentially in radio quiet objects and flat spectrum (compact) radio sources.

The origin of the outflows has also been subject of study. Correlations between the radio properties and the kinematic signatures of the outflows have suggested that the radio jets/lobes can play an important role to trigger such outflows both in radio loud and radio quiet objects (e.g. Humphrey et al. 2006; Leipsky & Bennert 2006; Heckman et al. 1984; Whittle et al. 1988, etc.). The dynamics of the ionized gas, which is subject to a partial or full radiation field from the central continuum source, may be affected by the radiation force as well (e.g. Boroson 2005). Finally, galactic superwinds may also exist. Star forming systems at low and high redshift are known to drive galactic winds with observable signatures in the wings of the emission lines (e.g. Shapiro et al. 2009).

Type 2 (obscured) quasars have been discovered in large quantities only in recent years (e.g. Zakamska et al. 2003; Martínez-Sansigre et al. 2005). From the observational point of view, these are the luminous counterparts of Seyfert 2 galaxies. In particular, Zakamska et al. identified in 2003 several hundred objects in the redshift range $0.3 \le z \le 0.8$ in the Sloan Digital Sky Survey (SDSS, York et al. 2000) with the high ionization narrow emission line spectra characteristic of type 2 AGNs and narrow line luminosities typical of type 1 quasars. Their far-infrared luminosities place them among the most luminous quasars at similar *z*. Several studies suggest that they have high star-forming luminosities (e.g. Zakamska et al. 2008; Hiner et al. 2009). They show a wide range of X-ray luminosities and obscuring column densities (Zakamska et al. 2004; Ptak et al. 2006). The host galaxies are most frequently ellipticals with irregular morphologies (Zakamska et al. 2006), and the nuclear optical emission is highly polarized in some cases (Zakamska et al. 2004; Lal & Ho 2010).

Because of the recent discovery, little is known about the possible existence of outflows in type 2 quasars. Greene et al. (2011) analysed the kinematics of the *extended* ionized gas in a sample of 15 luminous SDSS type 2 quasars at z < 0.5 in spatial scales of $\lesssim 10$ kpc. They found kinematically disturbed gas across the host galaxies and proposed that AGN driven outflows are stirring up the gas on scales of kpc.

We investigate in this paper the presence of ionized gas outflows in a sample of 13 optically selected type 2 quasars at $0.3 \le z \le$ 0.6. The observations and data analysis techniques are described in Section 2. The results are presented in Section 3 and discussed in Section 4. The results and conclusions are summarized in Section 5.

We assume $\Omega_{\Lambda} = 0.7$, $\Omega_{\rm M} = 0.3$, $H_0 = 71 \,\rm km \, s^{-1} \, \rm Mpc^{-1}$.

2 OBSERVATIONS AND ANALYSIS

The data were obtained with the Focal Reducer and Low Dispersion Spectrograph (FORS2) for the Very Large Telescope (VLT) installed on UT1 (Appenzeller et al. 1998). The observations were performed in two different runs: 2008 September 8 and 9 (four objects) and 2009 April 17 and 18 (nine objects). All spectra were obtained with the 600RI+19 grism and the GG435+81 order sorting filter. The useful spectral range was \sim 5030–8250 Å in the 2008 run and \sim 5300–8600 Å in the 2009 run, so that in all cases at least the H β and [O III] $\lambda\lambda$ 4959,5007 lines were within the observed spectral range. The pixel scales are $0.25 \text{ arcsec pixel}^{-1}$ and 0.83Å pixel⁻¹ in the spatial and spectral directions, respectively. The spectral resolution, as measured from the sky emission lines, was full width half-maximum (FWHM) = 7.2 ± 0.2 Å and 5.4 ± 0.2 Å for the 2008 and 2009 runs, respectively. The slit width was 1.3 arcsec in 2008 and 1.0 arcsec in 2009. The observations, data reduction and object sample are described in detail in Villar-Martín et al. (2010, 2011).

1-dim spectra were extracted from spatial apertures centered on the quasars spatial continuum centroid. The sizes were selected to optimize the S/N of the nuclear emission lines, in particular for the detection of faint broad wings. In general, the aperture sizes are \sim 1.5–2 arcsec, corresponding to physical sizes \sim 7–12 kpc depending on the object. Type 2 quasars are often associated with extended emission line regions (EELR) which can extend sometimes for tens of kpc (e.g. Villar-Martín et al. 2011), well beyond the size of the NLR. The apertures we have defined contain emission from both the NLR and, when existing, the EELR, although the contribution of the NLR is likely to dominate.

The spectral profiles of [O III] and $H\beta$ were fitted with the STAR-LINK DIPSO package. One or more Gaussian components were used depending on the quality of the fits. This was evaluated based both on the errors of the fits and a visual inspection of the residuals. The [O III] lines are in all cases the strongest and they were used

Table 1. Type 2 quasar sample studied in this work with kinematic measurements based on single Gaussian fits to the [O III] and H β lines. The redshift is shown in column (2). The [O III] and H β FWHM are are in columns (3) and (4). Column (5) gives the FWHM of the narrow [O III] core that remains after the asymmetric line wings are removed. AI10, AI20 and AI50 in columns (6), (7) and (8) quantify the degree of asymmetry of the [O III] line profile at 10, 20 and 50 per cent intensity levels, respectively. Positive, negative and zero values (taking errors into account) correspond to blue, red and no asymmetries, which are highlighted in bold, italic and normal fonts, respectively.

| Object | Z | FWHM [Ош] | FWHM H <i>β</i> | FWHM _{core} | AI10 | AI20 | AI50 |
|---------------|-------|---------------|--------------------|----------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| (1) | (2) | (3) | (4) | (5) | (6) | (7) | (8) |
| SDSS J2358-00 | 0.402 | 440 ± 50 | 420 ± 85 | 430 ± 10 | 0.03 ± 0.04 | 0.00 ± 0.03 | 0.00 ± 0.03 |
| SDSS J0025-10 | 0.303 | 435 ± 55 | 420 ± 70 | 320 ± 80 | $\textbf{0.33} \pm \textbf{0.05}$ | $\textbf{0.36} \pm \textbf{0.08}$ | $\textbf{0.42} \pm \textbf{0.09}$ |
| SDSS J0123+00 | 0.399 | 445 ± 45 | 470 ± 60 | 430 ± 50 | $\textbf{0.10} \pm \textbf{0.05}$ | 0.07 ± 0.07 | 0.04 ± 0.06 |
| SDSS J0217-00 | 0.344 | 910 ± 30 | 700 ± 60 | 700 ± 60 | $\textbf{0.22} \pm \textbf{0.03}$ | $\textbf{0.06} \pm \textbf{0.04}$ | -0.17 ± 0.09 |
| SDSS J0234-07 | 0.310 | 510 ± 60 | 420 ± 60 | 450 ± 50 | $\textbf{0.39} \pm \textbf{0.04}$ | $\textbf{0.52} \pm \textbf{0.05}$ | $\textbf{0.53} \pm \textbf{0.09}$ |
| SDSS J0955+03 | 0.422 | 1320 ± 30 | N/A | 940 ± 60 | $\textbf{0.13} \pm \textbf{0.03}$ | $\textbf{0.14} \pm \textbf{0.07}$ | -0.15 ± 0.09 |
| SDSS J1153+03 | 0.575 | 550 ± 10 | N/A | 440 ± 20 | -0.05 ± 0.06 | -0.05 ± 0.04 | -0.06 ± 0.09 |
| SDSS J1228+00 | 0.575 | 1860 ± 20 | N/A | 1100 ± 100 | -0.66 ± 0.03 | -0.41 ± 0.04 | -0.49 ± 0.05 |
| SDSS J1307-02 | 0.455 | 410 ± 10 | 455 ± 15 | 360 ± 10 | $\textbf{0.15} \pm \textbf{0.03}$ | -0.10 ± 0.05 | 0.00 ± 0.04 |
| SDSS J1337-01 | 0.329 | 900 ± 30 | N/A | 540 ± 50 | $\textbf{0.33} \pm \textbf{0.07}$ | $\textbf{0.19} \pm \textbf{0.07}$ | $\textbf{0.15} \pm \textbf{0.06}$ |
| SDSS J1407+02 | 0.309 | 430 ± 20 | 460 ± 15 | 400 ± 10 | $\textbf{0.18} \pm \textbf{0.06}$ | $\textbf{0.17} \pm \textbf{0.09}$ | $\textbf{0.20} \pm \textbf{0.10}$ |
| SDSS J1413-01 | 0.380 | 550 ± 10 | 570 ± 40 | 500 ± 10 | 0.03 ± 0.04 | 0.03 ± 0.03 | 0.00 ± 0.03 |
| SDSS J1546-01 | 0.383 | 500 ± 40 | 440 ± 30 | 430 ± 20 | $\textbf{0.39} \pm \textbf{0.08}$ | $\textbf{0.43} \pm \textbf{0.08}$ | 0.00 ± 0.05 |
| Median | 0.383 | 510 | 455 | 440 | 0.13 | 0.07 | 0.00 |
| STD | 0.091 | 440 | 93 | 230 | 0.28 | 0.24 | 0.26 |

to constrain the expected kinematic substructure of H β . The ratio between the [O III] $\lambda\lambda$ 4959,5007 components was forced to be 1:3, in accord with the theoretical value (Osterbrock 1989). Both components were forced to have the same FWHM and in general the separation in wavelength was also fixed. We found necessary to apply these constraints to all kinematic components in order to reject fits that were successful at reproducing the line shapes, but did not have physical meaning.

To start, we attempted to fit the [O III] profiles with a single Gaussian function, but this failed in all cases and very large residuals remained. The number of Gaussians was progressively increased until a good fit was achieved. Two or three Gaussians at most were required in all quasars. The H β line profile could be decomposed in different kinematic components only in some objects where the line had enough signal and was not severely affected by atmospheric or galactic absorption. In those cases, it was assumed that the line consists of the same number of kinematic components as [O III]. The FWHM and z of each component were constrained from those derived from [O III]. Only those components with a detectable flux were considered in the final fit of $H\beta$. Flux upper limits for the non-detected components were also estimated. The FWHM (corrected for instrumental broadening in quadrature), the velocity shift relative to the narrow [O III] core (see Section 2.1), the flux, the flux relative to the total line flux and the $[O_{III}]/H\beta$ ratio (when possible) were measured for all kinematics components.

The kinematic measurements obtained for the first five objects in Tables 1 and 2 are affected by slit effects (see Villar-Martín et al. 2011). The errors quoted take the resulting uncertainties into account.

The results of the fits are shown in Table 2 and Figs 1 to 13.

2.1 The systemic velocity

In order to characterize accurately the gas motions relative to their environment it is essential to determine the systemic redshift z_{sys} of the galaxy. The safest way to do this is to measure the stellar redshift. This is often not possible as for example in studies of type 1 AGNs, high-redshift galaxies, etc. where the stellar continuum cannot be isolated from the bright AGN contribution and/or it is not detected with enough signal. In such cases, alternative methods have been applied. Under the assumption that the BLR gas follows gravitational motions, the broad H β line centroid has been used to determine z_{sys} . For type 2 objects, the [O III] line fitted with a single Gaussian has often been used instead, both to measure z_{sys} and the stellar velocity dispersion. However, as different works have discussed (e.g. Boroson 2003, 2005; Greene et al. 2011, 2009; Husemann 2011), this is risky, since the NLR kinematics often reveal non-gravitational motions (see Section 1). Indeed, the very asymmetric and sometimes broad [O III] spectral profiles often make the determination of its own *z* uncertain by up to 200 km s⁻¹ (e.g. Heckman et al. 1981).

There are three objects in our sample with enough signal in the continuum and detected stellar features to try to estimate z_{sys} : SDSS J0217–00, SDSS J1337–01 (both with strong stellar features) and SDSS J0123+00 (faint features). For this, we use the full spectral fitting method with the starlink code (Cid-Fernandes et al. 2005), several sets of evolutionary stellar models (González Delgado et al. 2005; Charlot & Bruzual (in preparation) and two of the most up to date high/intermediate spectral resolution stellar libraries (Martins et al. 2005; Sánchez Blázquez et al. 2006).

We find that in all three cases there is a variety of models that produce acceptable fits (see one example in Fig. 14). These models allow a range of stellar redshifts which depend mainly on the single stellar population models used as templates for the fits and the weight given to the *G* band and Cau K stellar absorption features. The allowed redshifts imply a velocity range of ~100 km s⁻¹, which is the main uncertainty affecting the method.

Although the [O III] line in the nuclear spectra of our sample is in general quite broad and shows complex kinematic substructure (see Section 3), it shows a very narrow core whose redshift can be estimated with an accuracy of several km s⁻¹. We will use this as a measurement of z_{sys} .

Table 2. Results of the spectral decomposition analysis of [O III] and $H\beta$. (1) Object (shortened) name. (2) Sign of the [O III] profile asymmetry implied by AI10 (Table 1). (3) Flux in units of $\times 10^{-16}$ erg s⁻¹ cm⁻². (4) Fractional contribution of a given kinematic component to the total [O III] flux. (5) Idem for $H\beta$. (6) Ionization level as measured from the $[O III]/H\beta$ ratio. (7) FWHM. (8) Velocity shift V_S relative to the [O III] narrow core. (9) $[O III]/H\beta$ calculated with the total line fluxes (rather than the individual kinematic components as in 6). (10) Total [O III] luminosity in units of 10^{42} erg s⁻¹. (11) Stellar velocity dispersion calculated according to FWHM $[O III]_{core} = (1.24 \pm 0.76) \times FWHM_{stars}$ (Greene & Ho 2005). The errors do not account for the scatter of the correlation (see Section 3.2).

| Object | Asymm | Flux[O III] $\times 10^{-16}$ | $\frac{[O m]_i}{[O m]_{tot}}$ | $\frac{\mathrm{H}\beta_i}{\mathrm{H}\beta_{\mathrm{tot}}}$ | $\frac{[\text{O III}]_i}{\text{H}\beta_i}$ | FWHM[OIII] | $V_{\rm S}$ | $\frac{[O III]_{tot}}{H\beta_{tot}}$ | $L[O III]_{tot}$ $\times 10^{42}$ | FWHM _{stars} |
|---------|-------|--|--|--|--|---|--|--------------------------------------|--------------------------------------|----------------------------|
| (1) | (2) | erg s ⁻¹ cm ⁻² (3) | (4) | (5) | (6) | km s ⁻¹ (7) | km s ⁻¹ (8) | (9) | erg s ⁻¹ (10) | km s ⁻¹ (11) |
| 2358-00 | No | $\begin{array}{c} 75\pm 6\\ 50\pm 2\end{array}$ | $\begin{array}{c} 0.60 \pm 0.06 \\ 0.40 \pm 0.03 \end{array}$ | $\begin{array}{c} 0.62 \pm 0.08 \\ 0.38 \pm 0.05 \end{array}$ | $\begin{array}{c} 10\pm1.5\\ 11\pm1 \end{array}$ | $\begin{array}{c} 310\pm40\\ 1020\pm30 \end{array}$ | $\begin{array}{c} 10\pm40\\ 10\pm40 \end{array}$ | 10.9 ± 0.4 | 8.06 | 350 ± 40 |
| 0025-10 | Blue | 8.3 ± 0.4 35 ± 1 8.8 ± 0.9 | $\begin{array}{c} 0.16 \pm 0.01 \\ 0.67 \pm 0.03 \\ 0.17 \pm 0.02 \end{array}$ | $\leq 0.08 \ \sim 1 \ < 0.17$ | ≥ 9 3.4 ± 0.1 >4.3 | ≤ 130 500 ± 40 1330 ± 100 | $20 \pm 40 \\ -60 \pm 40 \\ -80 \pm 45$ | 4.6 ± 0.2 | 2.07 | 260 ± 60 |
| 0123+00 | Blue | 5.9 ± 0.9 61.7 ± 0.6 8.2 ± 0.7 | $\begin{array}{c} 0.08 \pm 0.01 \\ 0.81 \pm 0.02 \\ 0.11 \pm 0.01 \end{array}$ | <0.08 ~1 ≤0.13 | >8.3 8.8 ± 0.2 >7 | $\leq 200 \\ 470 \pm 40 \\ 1500 \pm 130$ | $20 \pm 50 \\ -15 \pm 50 \\ 3 \pm 50$ | 10.3 ± 0.3 | 5.21 | 350 ± 40 |
| 0217-00 | Blue | $\begin{array}{c} 30\pm2\\ 36\pm2 \end{array}$ | $\begin{array}{c} 0.46 \pm 0.03 \\ 0.54 \pm 0.03 \end{array}$ | $\begin{array}{c} 0.63 \pm 0.09 \\ 0.37 \pm 01.5 \end{array}$ | $\begin{array}{c} 10\pm2\\ 21\pm9 \end{array}$ | $\begin{array}{c} 470\pm50\\ 1530\pm40\end{array}$ | $\begin{array}{c} 80\pm45\\-120\pm50\end{array}$ | 14.6 ± 2.3 | 2.17 | 570 ± 50 |
| 0234-07 | Blue | $\begin{array}{c} 32.5\pm0.9\\ 32\pm1 \end{array}$ | $\begin{array}{c} 0.48 \pm 0.02 \\ 0.48 \pm 0.06 \end{array}$ | $\begin{array}{c} 0.50 \pm 0.04 \\ 0.40 \pm 0.06 \end{array}$ | $\begin{array}{c} 11.1\pm0.9\\ 14\pm2 \end{array}$ | $\begin{array}{c} 310\pm70\\ 950\pm30\end{array}$ | $\begin{array}{c} 0\pm 50\\ -270\pm 50\end{array}$ | 12.6 ± 0.6 | 2.77 | 360 ± 40 |
| 0955+03 | Blue | 0.9 ± 0.2 3.1 ± 0.4 2.6 ± 0.3 | $\begin{array}{c} 0.13 \pm 0.04 \\ 0.47 \pm 0.08 \\ 0.40 \pm 0.06 \end{array}$ | Unk. Unk. Unk. | Unk. Unk. Unk. | 470 ± 80 1110 ± 80 2500 ± 200 | 0 ± 20 190 ± 50 -620 ± 140 | 16 ± 4 | 1.54 | 740 ± 50 |
| 1153+03 | No | $\begin{array}{c} 26\pm1\\ 58\pm1\\ 9\pm2 \end{array}$ | $\begin{array}{c} 0.28 \pm 0.01 \\ 0.62 \pm 0.02 \\ 0.10 \pm 0.01 \end{array}$ | Unk. Unk. Unk. | Unk. 14 ± 4 Unk. | 270 ± 10 730 ± 40 1450 ± 80 | $5 \pm 10 \\ -5 \pm 10 \\ 330 \pm 70$ | 14.1 ± 0.1 | 15.7 | 360 ± 20 |
| 1228+00 | Red | 6.4 ± 0.3 4.6 ± 0.5 22.2 ± 0.4 | $\begin{array}{c} 0.19 \pm 0.01 \\ 0.14 \pm 0.2 \\ 0.67 \pm 0.02 \end{array}$ | Unk. Unk. Unk. | Unk. Unk. Unk. | 550 ± 20 780 ± 60 2300 ± 70 | -60 ± 10 640 ± 30 770 ± 20 | 12 ± 1 | 7.36 | 890 ± 80 |
| 1307-02 | Blue | $\begin{array}{c} 17\pm2\\ 19\pm2\\ 13.6\pm0.7\end{array}$ | $\begin{array}{c} 0.34 \pm 0.04 \\ 0.40 \pm 0.04 \\ 0.26 \pm 0.02 \end{array}$ | $\begin{array}{c} 0.23 \pm 0.03 \\ 0.61 \pm 0.05 \\ 0.16 \pm 0.04 \end{array}$ | 16 ± 2 7.0 ± 0.8 18 ± 4 | 250 ± 10 480 ± 20 1000 ± 30 | -13 ± 5 56 ± 9 -190 ± 20 | 10.9 ± 0.3 | 3.21 | 290 ± 8 |
| 1337-01 | Blue | 9.5 ± 0.4 22.3 ± 0.5 | $\begin{array}{c} 0.30 \pm 0.01 \\ 0.70 \pm 0.02 \end{array}$ | Unk. Unk. | Unk. Unk. | $\begin{array}{c} 350\pm20\\ 1280\pm60 \end{array}$ | $\begin{array}{c} 30\pm 6\\ -290\pm 10\end{array}$ | $25 \pm 2^*$ | 2.03 | 440 ± 40 |
| 1407+02 | Blue | 55 ± 1 41 ± 2 15.0 ± 0.6 | $\begin{array}{c} 0.50 \pm 0.02 \\ 0.37 \pm 0.02 \\ 0.13 \pm 0.01 \end{array}$ | $\begin{array}{c} 0.57 \pm 0.03 \\ 0.43 \pm 0.04 \\ < 0.18 \end{array}$ | 9.4 ± 0.5 9.0 ± 0.9 >6 | 290 ± 10 720 ± 20 1670 ± 120 | 10 ± 3 -100 \pm 10 -70 \pm 30 | 10.9 ± 0.4 | 2.03 | 320 ± 8 |
| 1413-01 | No | 14.2 ± 2 45 ± 2 14 ± 2 | $\begin{array}{c} 0.20 \pm 0.02 \\ 0.61 \pm 0.03 \\ 0.19 \pm 0.02 \end{array}$ | $\begin{array}{c} 0.19 \pm 0.02 \\ 0.56 \pm 0.04 \\ 0.25 \pm 0.03 \end{array}$ | 9 ± 1 10.0 ± 0.6 7 ± 1 | 260 ± 20 600 ± 20 1190 ± 50 | -20 ± 10 40 ± 8 -145 ± 30 | 9.6 ± 0.4 | 3.07 | 400 ± 8 |
| 1546-01 | Blue | 7.6 ± 0.1 5.5 ± 0.5 | $0.58 \pm 0.02 \\ 0.42 \pm 0.04$ | $0.71 \pm 0.08 \\ 0.29 \pm 0.08$ | $\begin{array}{c} 6.5\pm0.7\\ 12\pm3.5\end{array}$ | $\begin{array}{c} 340\pm20\\ 780\pm30\end{array}$ | $\begin{array}{c} 00\pm20\\ -270\pm35 \end{array}$ | 8.3 ± 0.3 | 5.21 | 350 ± 20 |

In the three objects mentioned above, it is found that this core is at $\lesssim 100 \,\mathrm{km \, s^{-1}}$ of the stellar velocity, the uncertainty being due to the range of stellar redshifts allowed by the models. Given the difficulty to infer $z_{\rm sys}$ from the stars in most quasar spectra studied here, we will use the narrow [O III] core in all objects instead. This method is supported by the results of Greene & Ho (2005). They compared the gaseous and stellar kinematics in the nuclei of a large sample of active galaxies using spectra of somewhat better resolution ($R = \lambda/\Delta\lambda \sim 1800$ versus $R \sim 900-1400$ at the [O III] wavelength in our spectra, depending on the object). They found that the FWHM of the [O III] core, i.e. after the asymmetric wings were removed, and FWHM_{stars} are correlated (although with a large scatter) according to FWHM_{core} = (1.24 ± 0.76) × FWHM_{stars}.

Based on this result, they propose that $FWHM_{core}$ is a reasonable tracer of the stellar velocity dispersion. It seems reasonable to expect that it also provides a good estimation of the stellar redshift and therefore z_{sys} .

Greene et al. (2009) questioned later the validity of this result for luminous obscured AGNs, based on the lack of correlation between the $[O_{III}]$ FWHM and the stellar velocity dispersion. However, the analysis is based on single Gaussian fits to the emission lines that ignore their complex kinematic substructure. Given the high complexity of the line profiles (see Section 3.2), it is not surprising that no correlation is found. As Table 1 shows, the FWHM of the core is often noticeably smaller than that resulting from a single Gaussian fit to the line profile.



Figure 1. Fit to H β and [O III] $\lambda\lambda$ 4959,5007 for SDSS J2358–00. In each figure from 1 to 13 the data and the fit are shown in black and orange, respectively. The same colour and line style are used for each kinematic component in all emission lines ([O III] $\lambda\lambda$ 4959,5007 and, when possible, also H β). Red and blue are used for the most redshifted and blueshifted components, respectively. Green is used for an intermediate redshift. When two components are identified with equal *z* within the errors (as the example in this figure), the same colour but different line style are used for each one. The vertical black line marks the wavelength of the very narrow [O III] peak, which we have used to calculate the velocity shifts.



Figure 2. SDSS J0025-10. Line styles and colours as in Fig. 1.

2.2 The identification of kinematically perturbed gas

We are specially interested in those gas components whose kinematic properties suggest non-gravitational, perturbed motions. In order to identify them, we need to compare the FWHM of each kinematic component with the stellar velocity dispersion FWHM_{stars}. Similar values could be explained if the gas is supported by random motions,¹ while rotation would produce narrower FWHM for the gas than for the stars. Broader gas components would imply kinematic perturbation.

In general, it is not possible to measure $FWHM_{stars}$ directly from our spectra. Instead, following Greene & Ho (2005) we will infer the value for each object using $FWHM_{core}$ (Section 2.1). Although the scatter of the correlation between these two parameters pro-



Figure 3. SDSS J0123+00. Line styles and colours as in Fig. 1.



Figure 4. SDSS J0217–00. Line styles and colours as in Fig. 1.



Figure 5. SDSS J023–07. Line styles and colours as in Fig. 1.

duces large uncertainties, the FWHM_{core} values in Table 1 suggest FWHM_{stars} in the range \sim 250–900 km s⁻¹ (Table 2), consistent with values measured in other type 2 quasars (Bian 2007; Liu et al. 2009) using stellar features.

¹ Type 2 quasars are most frequently associated with elliptical galaxies (Zakamska et al. 2006; Marinieri et al. 2011), so random motions are possible.



Figure 6. SDSS J0955+03. Line styles and colours as in Fig. 1.



Figure 7. SDSS J1153+03. Line styles and colours as in Fig. 1.



Figure 8. SDSS J1228+00. Line styles and colours as in Fig. 1.



Figure 9. SDSS J1307–02. Line styles and colours as in Fig. 1.



Figure 10. SDSS J1337–01. Line styles and colours as in Fig. 1.



Figure 11. SDSS J1407+02. Line styles and colours as in Fig. 1.



Figure 12. SDSS J1413–01. Line styles and colours as in Fig. 1.



Figure 13. SDSS J1546–00. Line styles and colours as in Fig. 1.

The velocity shifts $V_{\rm S}$ of the different [O III] kinematic components relative to the narrow line core will provide additional information to characterize the kinematics.

3 RESULTS

3.1 Integrated line profiles

We show in Table 1 the FWHM of [O III] and $H\beta$ lines. For an initial, general characterization of these quantities we have applied single Gaussian fits to both lines. The goal of this is to compare with similar studies of type 1 quasars.

The [O III] FWHM values in Heckman et al. (1984) subsample of 35 type 1 quasars are in the range [260–850] km s⁻¹, with a median value of 460 km s⁻¹. If only the 10 radio quiet quasars are considered the range is the same, but the median value increases to 570 km s^{-1} . The range in our sample is [410 ± 10 to 1860 ± 20] km s⁻¹ and the median is 510 km s^{-1} .

A visual inspection of the [O III] spectral profiles reveals that it is asymmetric in the majority of objects. Following Heckman et al. (1981), we have quantified the degree of asymmetry at 10, 20 and 50 per cent of the line maximum intensity using the parameters AI10, AI20 and AI50. AI10 is defined as (WL10 - WR10)/(WL10 +WR10), where WL10 and WR10 are the half widths to the left and right of the centre of the narrow [O III] core at 10 per cent intensity



Figure 14. Fit to the continuum of SDSS J0217–00 using stellar population synthesis models. The spectrum has been translated to z = 0. The fluxes have been normalized to the flux at 4020 Å. Different templates produce a good fit, so that a range of stellar (and therefore systemic) redshifts is allowed. These imply a velocity range of ~100 km s⁻¹.



Figure 15. Schematic representations of the various parameters used to quantify the degree of asymmetry of the $[O \ m]$ line profile at different intensity levels, following Heckman et al. (1981).

level. The widths were corrected for instrumental broadening. AI20 and AI50 are defined in the same way, but at 20 and 50 per cent intensity levels, respectively. AI10 is introduced compared with previous studies (see also Spoon & Holt 2010) because we find that in most objects the asymmetry is perceived at very low intensity levels, in the form of faint wings. The schematic representation of the various parameters involved in these calculations is shown in Fig. 15. The half widths WL10', WR10', etc have a ' ' ' because they correspond to values not corrected for instrumental broadening.

AI10, AI20 and AI50 are shown in Table 1. Positive, negative and zero values indicate blue (highlighted in bold), red (italic fonts) and no asymmetries (normal fonts), respectively. It can be seen that the fraction of asymmetric profiles increases at lower intensity levels (7/13 at 50 per cent; 9/13 at 20 per cent and 10/13 at 10 per cent levels). While at 50 per cent level there is a variety of asymmetries with no clear preference for a given sign, at lower intensity blue asymmetries become predominant: at 10 per cent level 9/13 objects show a blue asymmetry, versus 2 with red and 2 more which no asymmetry.

Our results are consistent with Heckman et al. (1984), who found a clear predominance of blue asymmetric [O III] profiles in radio quiet and flat steep spectrum (compact) radio loud quasars at $z \le 0.5$. The asymmetries disappear in steep spectrum (extended) radio loud objects. None of the objects in our sample are radio loud.

An excess of blue asymmetries suggests that a significant fraction of the line emitting gas is moving radially in an outflow (see Section 1).

3.2 Spectral decomposition of the H β -[O III] line profiles.

The [O III] spectral line profiles show complex kinematic substructure in all 13 objects: three kinematic components are required to produce the best fit in 8/13 objects, while two are required for the remaining five (Figs 1–13).

All quasars in our sample have one or two kinematic components with [O III] FWHM \leq FWHM_{stars} (Table 2), i.e. they show no sign of kinematic disturbance. The presence of two components which seem to follow gravitational motions is not surprising, since the emissions from the NLR (probably dominant) and possibly also the EELR are expected within the spatial apertures used to extract the spectra. According to studies of narrow line radio galaxies, the EELR is expected to have different kinematic, physical and ionization properties than the NLR (e.g. Tadhunter, Fosbury & Quinn 1989; Robinson et al. 1987).

All objects show, moreover, a very broad component with [O III] FWHM $\geq 2 \times$ FWHM_{stars}. In general, this component is broader than the maximum possible value of FWHM_{stars} allowed by Greene & Ho (2005) correlation scatter (see Section 2.2.). Therefore, the uncertainty on FWHM_{stars} does not affect our conclusion. Thus, the NLR of all type 2 quasars in our sample contains highly perturbed gas, in addition to the more quiescent, ambient gas whose FWHM is consistent with gravitational motions.

The FWHM of the perturbed gas components are in the range 780 ± 30 to 2500 ± 200 km s⁻¹ with a median value of 1280 km s⁻¹. Their velocity shifts relative to the [O III] narrow core V_S are in the range 10 ± 40 and up to $\sim -620 \pm 140$ km s⁻¹, with a median value of ~ -145 km s⁻¹ (see also Greene et al. 2011).

We have plotted $R = FWHM[O III]/FWHM_{stars}$ versus V_S for all kinematic components (Fig. 16). The two objects highlighted in colour are SDSS J1153+03 (orange) and SDSS J1228+00 (green). Villar-Martín et al. (2011) proposed that these quasars might have an intermediate orientation between type 1 and type 2, so that we have a direct view of an intermediate density region, more interior and closer to the black hole than the NLR we see in the rest of the sample. Fig. 16 shows that they are also peculiar from the kinematic point of view, since they are the only ones showing kinematic components with large red velocity shifts.

If these two peculiar objects are excluded, Fig. 16 shows a strong trend for the highly perturbed gas (i.e. kinematic components with $R \gtrsim 2$) to be *blueshifted*, while the more quiescent gas (components with $R \lesssim 1.0$) shows in general no or small velocity shifts.

 $V_{\rm S}$ is plotted versus the FWHM in Fig. 17 for the individual kinematic components isolated in each quasar (the two peculiar quasars are excluded). The same symbol is used for each object. Selecting a given symbol, it can be seen that the largest FWHM values (i.e. the most perturbed gas in a given object) are in general associated with the largest blueshifts (see also Table 2). This result is found in all objects where the velocity uncertainties allowed a definitive conclusion (7/11).

Thus, the perturbed gas shows both broad FWHM compared with the stellar velocity dispersion and a trend to show the largest



Figure 16. Velocity shift *V*s in km s⁻¹ versus $R = \text{FWHM}[\text{O III}]/\text{FWHM}_{\text{stars}}$ for all kinematic components isolated in the spectral fits. Coloured symbols correspond to the two peculiar objects suspected to have intermediate orientation between type 1 and 2 (green for SDSS J1228+00; orange for SDSS J1153+03). Excluding these, there is a global trend for the perturbed gas components ($R \gtrsim 2$) to be bluesfhited, while more quiescent kinematic components $R \lesssim 1$) show small or null velocity shifts.



Figure 17. Velocity shift *V*s (km s⁻¹) versus the FWHM (km s⁻¹) in log of all kinematic components isolated in the the [O III] line for all quasars in the sample, except the two peculiar objects SDSS J1228+00 and SDSS J1153+03. Equal symbols correspond to the same object. There is a general trend such that in each object the broadest component shows also the largest blueshift.

blueshift. The perturbed gas is therefore responsible for the blue line profile asymmetries. All together suggest the presence of ionized gas outflows in the inner \lesssim few kpc, such that the receiding gas is obscured by dust.

The relative contribution of the perturbed (i.e. outflowing) gas to the total [O III] flux $R2 = [O III]^{outf}/[O III]^{tot}$ (column 5 in Table 2) varies strongly from object to object. It can account for ~10 per cent and up to ~70 per cent of the total line flux. R2 is plotted versus Rin Fig. 18. An anticorrelation is found such that, the more perturbed the outflowing gas is (larger R), the lower its relative contribution to the total [O III] flux. A Spearman's rank correlation coefficient $\rho =$ -0.79 is found and a statistical significance at the 99.95 per cent level.

The perturbed gas is also detected in H β in seven objects. The [O III]/H β ratio is in general very high (\geq 10, in most cases where it could be measured), which is similar or even larger than that of the narrower kinematic components. This suggests very high excitation level of the outflowing gas.



Figure 18. Relative contribution *R*2 to the total [O III] line flux of the perturbed component (column 4 in Table 2) versus $R = FWHM/FWHM_{stars}$ for the perturbed kinematics components (i.e. the outflowing gas). An anticorrelation is observed. Kinematic components with more extreme kinematics (larger *R*) have a smaller relative contribution to the total line flux.

Given the size of the apertures used to extract the spectra, the effects of the outflows we see are constrained within radial spatial scales of $r \leq$ few kpc from the central engine. Larger spatial scales of the total outflowing region are not discarded (e.g. Humphrey et al. 2010).

4 DISCUSSION

4.1 Outflows in the NLR of type 2 quasars

We have found evidence for outflows of ionized gas in the central regions ($r \leq$ few kpc) of most type 2 quasars in our sample (11/13). Six objects in the sample were selected on the basis (among other things) of prior evidence of nuclear kinematic perturbation (Villar-Martín et al. 2011) such as broad and/or highly asymmetric [O III] lines. However, not obvious bias was applied to give preference to blue versus red asymmetries. Moreover, six out of the seven objects for which kinematic criteria were not applied also contain outflows. So, in spite of the apparent selection bias, the results suggest that the outflow phenomenon is common in the NLR of type 2 quasars. This is consistent with Heckman et al. (1984), who found similar evidence in the NLR of the majority of radio quiet type 1 quasars (see also Leipsky & Bennert 2006).

The outflowing gas shows high ionization level with $[O \text{ III}]/\text{H}\beta$ sometimes $\gtrsim 10$, and often similar or larger than the more quiescent gas. A similar result has been found in some radio quiet type 1 quasars (e.g. Leipsky & Bennert 2006) and Seyfert galaxies (Veilleux 1991). Independently of the outflow origin, such high excitation suggests that stars are not responsible for ionizing the outflowing gas, but rather AGN related phenomena. This supports that the outflows are within the quasar ionization cones, as commented above. Although contribution from the EELR is expected in the spectra of at least some objects, the line emission is likely to be dominated by the NLR so that the outflows are expected to be located in this region.

The outflowing gas emits very broad lines (FWHM $\gtrsim 2 \times$ FWHM_{stars}, with FWHM in the range 780 \pm 30 to 2500 \pm 200 km s⁻¹) and *projected* shifts relative to the systemic velocity in the range $V_{\rm S} \sim 10 \pm 40$ and up to $\sim -620 \pm 140$ km s⁻¹ relative to the systemic velocity. For comparison, Nesvadba et al. (2011) measured FWHM of up to ~ 5000 km s⁻¹ and $V_{\rm S} \sim -1550$ km

 s^{-1} for the outflowing gas in a type 2 quasar at z = 3.5. The V_S are in general much less extreme than the motions implied by the large FWHM. Very broad lines can be produced by acceleration of clouds by or behind the bow shock generated by the outflow (e.g. Villar-Martín et al. 1999). Also, it must be taken into account that the deprojected velocity shifts are likely to be larger, since these are type 2 objects and the cones axis is close to the plane of the sky.

We have found a anticorrelation such that the strongest the kinematic perturbation (larger R), the lowest the relative contribution of the outfowing gas to the total [O III] line (smaller R2). Since there is no obvious change in the ionization level of the outfowing gas relative to the ambient, non perturbed gas, this anticorrelation possibly reflects that, the more perturbed the gas is, the smaller the gas mass fraction the outflow involves.

Based on the spatially resolved behaviour of the FWHM and $V_{\rm S}$ of [O III] line in a sample of 15 type 2 quasars at 0.11 $\leq z \leq$ 0.43, Greene et al. (2011) also proposed that outflows of ionized gas are present in their sample.

4.2 Outflow mass and mass outflow rate

The mass of the ionized outflowing gas M can be estimated with

$$M = \frac{1.4m_p L(\mathrm{H}\beta)}{n_e h v_{\mathrm{H}\beta} \alpha_{\mathrm{H}\beta}^{\mathrm{eff}}}$$

Ì

where L(H β) is the luminosity of the broad H β component, n_e the density of the outflowing gas, m_p the proton mass, $\alpha_{H\beta}^{eff}$ the effective H recombination coefficient for H β (Osterbrock 1989) and $h\nu_{H\beta}$ is the energy of an H β photon.

This refers to the outflowing gas detected within the slit. If the outflows are constrained to a region of size r <few kpc, we expect M to provide an adequate measurement of the total outflowing gas, otherwise (e.g. Humphrey et al. 2010) the inferred masses are lower limits.

Large uncertainties affect this calculation since the unreddened H β luminosity is unknown and we have no density measurements. The spectral fits imply observed L(H β) in the range (0.2–2.1) × 10⁴¹ erg s⁻¹.

Using the [S II] doublet from the SDSS spectra of type 2 quasars, Greene et al. (2011) find a range of densities \sim 250–500 cm⁻³, with a mean value of 335 cm^{-3} . [O II] $\lambda 3727$ is within the spectral range for SDSS J0217-00. The outflow is also evident in this line, which presents a clear blue asymmetry. Given the clear detection of this wing, it is reasonable to think that *n* is below the critical density of this line $n_{\rm crit} = 3 \times 10^3 \,{\rm cm}^{-3}$. The gas densities estimated in other objects (radio galaxies, Seyferts, far-infrared galaxies with galactic superwinds) for the outflowing (nuclear or extended) gas cover a broad range from \sim several hundred cm⁻³ (e.g Heckman, Armus & Miley 1990; Arribas & Mediavilla 1993; Villar-Martín et al. 1999) to $n_e > 10^{4.5}$ (Axon et al. 1998). $n = 250 \,\mathrm{cm}^{-3}$ seems therefore a reasonable lower limit. Thus, we assume a range of densities n \sim 250–3000 cm⁻³ for the outflowing gas. The implied values for M using the uncorrected L(H β) are in the range \sim (0.3–1.5) \times 10⁷ M_{\odot} for $n \sim 250 \,\mathrm{cm}^{-3}$ and depending on the object. The values are 120 times lower for $n \sim 3000 \,\mathrm{cm^{-3}}$. Greene et al. (2011) propose $\sim 10^7 \text{ M}_{\odot}$ as a strict lower limit assuming a gas density $\sim \text{few} \times$ 100 cm⁻³. It must be taken into account that these authors argue that all the H β emitting gas is involved in the outflow (we find this is not the case), and that much lower densties are possible.

For an illustrative comparison, the masses involved in *neutral* and *molecular* outflows in Seyfert and ultraluminous infrared galaxies (ULIRGs), which are constrained to spatial scales of \ll few kpc, are

 $>10^8$ M_{\odot}, and often $>10^9$ M_{\odot} (e.g. Rupke, Veilleux & Sanders 2005; Sturm et al. 2011).

It is obvious that better constraints on the intrinsic L(H β) and density of the outflowing gas are essential to obtain more tightly constrained masses. It is also essential to investigate the existence of outflows in other gas phases. Indeed, outflow studies in different types of galaxies suggest that the ionized phase reveals a very small fraction of the total outflowing mass, which is likely to be dominated by the molecular and neutral phases. As an example, the mass of ionized gas in the (galactic) superwind region of the M82 starburst galaxy is $\sim 2 \times 10^5$ M_{\odot} (Heckman et al. 1990), while the molecular mass involved in the outflow is $> 3 \times 10^8$ M_{\odot} (Walter, Weiss & Scoville 2002).

The mass outflow rate \dot{M} can be calculated as (Dopita & Sutherland 2003)

$$\dot{M} = \frac{3v_o L(\mathrm{H}\beta)m_p}{rnhv_{\mathrm{H}\beta}\alpha_{\mathrm{H}\beta}^{\mathrm{eff}}} = \frac{2.1v_0 M}{r}.$$

Assuming a spherical geometry, v_0 is the expanding velocity of the outflowing bubble and *r* its radius. In addition to those affecting *M*, additional uncertainties are involved for v_0 , which is likely to be $> V_s$ (Table 2) and *r*. Although different works suggest that the radius of the NLR in quasars is ~1-several kpc (e.g. Bennert et al. 2002; Greene et al. 2011), this is only an upper limit for *r*, since the size of the outflowing region could be smaller.

In spite of the uncertainties, it is interesting to study which conditions are required to produce different \dot{M} . For instance, mass flow rates of the NLR outflows discovered in Seyfert and nearby narrow line radio galaxies are typically \leq few M_O yr⁻¹ (Heckman et al. 1981; Rupke et al. 2005; see also Humphrey et al. 2010). Such values are certainly possible in our sample within a rather broad range of allowed parameter values.

More restrictive conditions are required to produce $\dot{M} > 50 \,\mathrm{M}_{\odot} \,\mathrm{yr}^{-1}$ as observed in some galactic scale outflows powered by active star formation (e.g. Heckman, Armus & Miley 1990; Lehnert & Heckman 1996; Rupke, Veilleux & Sanders 2002). Let us consider a scenario which favours the efficiency of the outflow: low densities ($n_e = 250 \,\mathrm{cm}^{-3}$) and large line luminosity corrections (e.g. intrinsic $L(H\beta) \sim 10 \times \mathrm{observed} \, L(H\beta)$). With these favourable prerequisites, the size of the outflowing region must be $r \ll 1$ kpc. Studies of ionized gas outflows using ultraviolet and X-ray absorption lines indicate that such small distances relative to he continuum source are possible. If the outflow region has $r \sim \mathrm{several}$ kpc as argued by Greene et al. (2011), it is very unlikely that such high mass outflow rates are generated, since most of the outflowing gas should be collimated within a very narrow cone near the plane of the sky for *all* objects.

It is clearly essential to determine the location of the outflows to constrain the mass outflow rates.

5 CONCLUSIONS

We have analysed the spatially integrated kinematic properties of the NLR gas in a sample of 13 optically selected SDSS type 2 quasars at $z \sim 0.3$ –0.6 using the [O III] $\lambda\lambda$ 4959,5007 lines. Optical long-slit spectroscopy at moderate spectral resolution $R \sim 900$ – 1300 obtained with FORS2 on the VLT has been used for this purpose.

The line profiles show a significant asymmetry in 11 of the 13 quasars. There is a clear preference for blue asymmetries, which are found in 9/13 objects. An excess of blue line asymmetries have

been observed in other active classes and non active galaxies. They are suggestive of outflows, such that the receding expanding gas is preferentially obscured. We propose the same interpretation. The outflows we see are constrained within spatial scales of r < few kpc from the central engine.

This scenario is favoured by other results we find: in addition to more quiescent gas, whose kinematic properties are consistent with gravitational motions, we have discovered highly perturbed gas in all objects. This gas emits very broad lines ($R = FWHM[OIII]/FWHM_{stars} \gtrsim 2$). While the quiescent gas shows small or null velocity shifts relative to the systemic velocity, the highly perturbed gas trends to show larger shifts. These, moreover, are blueshifts in general. In each object the most perturbed gas trends to have the largest blueshift as well. All together support that the perturbed gas, which is responsible for the blue asymmetries on the line profiles, is outflowing.

The *projected* velocity shifts relative to the narrow core of the [O III] line, which we have considered a reasonable tracer of the systemic velocity, are in the range $\sim 10 \pm 40$ and up to $\sim -620 \pm 140 \text{ km s}^{-1}$, with a median value of $\sim -145 \text{ km s}^{-1}$. The intrinsic expanding velocities are likely to be larger, given the orientation of the objects. We FWHM for the outflowing gas in the range 780 ± 30 to $2500 \pm 200 \text{ km s}^{-1}$ with a median value of 1280 km s^{-1} . The relative contribution of the outflowing gas to the total [O III] line flux varies from object to object in the range $\sim 10-70$ per cent. An anticorrelation is found such that, the more perturbed the outflowing gas is (larger *R*), the lower its relative contribution is to the total [O III] flux. Our results suggest that outflows with more perturbed kinematics involve a smaller fraction of the total mass of ionized gas.

The outflowing gas is in general very highly ionized, as suggested by the large $[O \amalg]/H\beta$ values (often $\gtrsim 10$). Independently of the outflows origin, this suggests that the gas is ionized by the quasar, rather than stars, an therefore it is located within the quasar ionization cones.

We argue that our results suggest that nuclear dusty outflows are a common phenomenon in optically selected type 2 quasars at $0.3 \lesssim z \lesssim 0.6$.

ACKNOWLEDGMENTS

This work has been funded with support from the Spanish Ministerio de Ciencia e Innovación through the grants AYA2007-64712, AYA2009-13036-C02-01 and ESP2007-65475-C02-01 and co-financed with FEDER funds. Thanks to Clive Tadhunter and the staff at Paranal Observatory for their support during the observations. The Sloan Digital Sky Survey (SDSS) is a joint project funded by the Alfred P. Sloan Foundation, the SDSS member institutions, NASA, the National Science Foundation, the U.S. Department of Energy, and Monbusho. The official SDSS web site is www.sdss.org.

REFERENCES

- Appenzeller I. et al., 1998, Messenger, 94, 1
- Arribas S., Mediavilla E., 1993, ApJ, 410, 552
- Axon D., Marconi A., Capetti A., Maccetto F., Schreier E., Robinson A., 1998, ApJ, 496, L75
- Bennert N., Falcke H., Schulz H., Wilson A., Wills B., 2002, ApJ, 547, 105
- Bian W., 2007, in Ho L. C., Wang J. M., eds, ASP Conf. Ser. Vol. 373, The Central Engine of Active Galactic Nuclei. Astron. Soc. Pac., San Francisco, p. 675
- Bian W., Yuan Q., Zhao Y., 2005, MNRAS, 364, 187
- Boroson T., 2003, ApJ, 585, 647

- Boroson T., 2005, AJ, 130, 381
- Cid-Fernandes R., Mateus A., Sodr L., Stasińska G., Gomes J., 2005, MNRAS, 358, 363
- Colbert E., Baum S., Gallimore J., O'Dea C., Lehnert M., Tsvetanov Z., Mulchaey J., Caganoff S., 1996, ApJS, 105, 75
- Crenshaw D., Kraemer S., George I., 1999, ARA&A, 2003, 41, 117
- Crenshaw D., Schmitt H., Kraemer S., Mushotzky R., Dunn J., 2010, ApJ, 708, 419
- Di Matteo T., Springel V., Hernquist L., 2005, Nat, 433, 605
- Dopita M., Sutherland R., 2003, Astrophysics of the Diffuse Universe. Springer, Berlin, New York, Springer. ISBN 3540433627
- Ferrarese L., Merritt D., 2000, ApJ, 539, L9
- Gebhardt K., Bender R., Bower G., 2000, ApJ, 539, L13
- González Delgado R., Cerviño M., Martins L., Leitherer C., Hauschildt P., 2005, MNRAS, 357, 945
- Greene J., Ho L., 2005, ApJ, 630, 122
- Greene J., Zakamska N., Liu X., Barth A., Ho L., 2009, ApJ, 702, 441
- Greene J., Zakamska N., Ho L., Barth A., 2011, ApJ, 732, 9
- Heckman T., Miley G., van Breugel W., Butcher H., 1981, ApJ, 247, 403
- Heckman T., Miley G., Green R., 1984, ApJ, 281, 525
- Heckman T., Armus L., Miley G., 1990, ApJS, 74, 833
- Hiner K., Canalizo G., Lacy M., Sajina A., Armus L., Ridgway S., Storrie-Lombardi L., 2009, ApJ, 706, 508
- Holt J., Tadhunter C., Morganti R., 2008, 387, 639
- Humphrey A., Villar-Martín M., Fosbury R., Vernet J., di Serego Alighieri S., 2006, MNRAS, 369, 1103
- Humphrey A., Villar-Martín M., Sánchez S., Martínez-Sansigre A., González Delgado R., Pérez E., Tadhunter C., Pérez-Torres M., 2010, MNRAS, 408, L1
- Husemann B., 2011, PhD thesis, Univ. Potsdam
- Komossa S., Xu D., Zhou H., Storchi-Bergmann T., Binette L., 2008, ApJ, 680, 926
- Lal D. V., Ho. L., 2010, AJ, 139, 1089
- Lehnert M., Heckman T., 1996, ApJ, 462, 651
- Leipski C., Bennert N., 2006, A&A, 448, 165
- Liu X., Zakamska N., Greene J., Strauss M., Krolik J., Heckman T., 2009, ApJ, 702, 1098
- Mainieri V. et al., 2011, A&A, submitted (arXiv:1105.5395)
- Martínez-Sansigre A., Rawlings S., Lacy M., Fadda D., Marleau F., Simpson C., Willott C., Jarvis M., 2005, Nat, 436, 666

- Martins L. P., Gonzalez Delgado R. M., Leitherer C., Cervino M., Hauschildt P. H., 2005, MNRAS, 358, 49
- Nesvadba N., Polletta M., Lehnert M., Bergeron J., de Breuck C., Lagache G., Omont A., 2011, MNRAS, 415, 2359
- Osterbrock D. E., 1989, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei. University Science Books, Mill Valley, CA
- Ptak A., Zakamska N., Strauss M., Krolik J., Heckman T., Schneider D., Birnkmann J., 2006, ApJ, 637, 147
- Robinson A., Binette L., Fosbury R., Tadhunter C., 1987, MNRAS, 227, 97
- Rupke D., Veilleux S., Sanders D., 2002, ApJ, 570, 588
- Rupke D., Veilleux S., Sanders D., 2005, ApJ, 632, 751
- Shapiro K. et al., 2009, ApJ, 701, 955
- Spoon H., Holt J., 2009, ApJ, 702, 42
- Sturm E. et al., 2011, ApJ, 733, L16
- Sanchez-Blazquez P. et al., 2006, MNRAS, 371, 703
- Tadhunter C., Fosbury R., Quinn P., 1989, MNRAS, 240, 225
- Veilleux S, 1991, ApJ, 369, 331
- Villar-Martín M., Tadhunter C., Morganti R., Axon D., Koekemoer A., 1999, MNRAS, 307, 24
- Villar-Martín M., Tadhunter C., Humphrey A., Pérez E., Martínez-Sansigre A., González-Delgado R., Pérez-Torres M., 2010, MNRAS, 407, L6
- Villar-Martín M., Tadhunter C., Humphrey A. R., Fraga-Encinas R., González Delgado M., Pérez Torres A., Martínez-Sansigre A., 2011, MNRAS, 416, 262
- Walter F., Weiss A., Scoville N., 2002, AJ, 580, L21
- Whittle M., Pedlar A., Meurs E., Unger S., Axon D., Ward M., 1988, ApJ, 326, 125
- York D. G. et al., 2000, AJ, 120, 1579
- Zakamska N. et al., 2003, AJ, 126, 2125
- Zakamska N., Strauss M., Heckman T., Ivezić Z., Krolik J., 2004, AJ, 128, 1002
- Zakamska N. et al., 2005, AJ, 129, 1212
- Zakamska N., Strauss M., Krolik J. et al., 2006, AJ, 132, 1496
- Zakasmka N., Gómez L., Strauss M., Krolik J., 2008, ApJ, 136, 1607

This paper has been typeset from a TEX/LATEX file prepared by the author.